

Cohomological classification of Ann-functors

Nguyen Tien Quang and Dang Dinh Hanh

April 27, 2009

Abstract

Regular Ann-functor classification problem has been solved with Shukla cohomology. In this paper, we would like to present a solution to the above problem in the general case and in the case of strong Ann-functors with, respectively, Mac Lane cohomology and Hochschild cohomology.

Mathematics Subject Classifications (2000): 18D10, 13D03.

Key words: Ann-category, Ann-functor, classification, Mac Lane ring cohomology, Hochschild cohomology.

1 Introduction

The definition of *Ann-categories* was presented by N.T.Quang [7] in 1987, which is regarded as a categorization of ring structure. Each Ann-category $\mathcal{A}$ is Ann-equivalent to its reduced Ann-category. This Ann-category is of the type (R, M, h) , where R is a ring of congruence classes of objects of $\mathcal{A}$, $M = \text{Aut}(0)$ is the R -bimodule and h is a 3-cocycle in $Z_{MaL}^3(R, M)$ (due to Mac Lane [6]). Then, there exists a bijection between the set of congruence classes of Ann-categories of the type (R, M) and the cohomology group $H_{MaL}^3(R, M)$ (see [11]). For the *regular Ann-categories* (whose the commutativity constraint satisfies $c_{X,X} = id$), then in the above bijection, the group $H_{MaL}^3(R, M)$ is replaced with the Shukla cohomology group $H_{Sh}^3(R, M)$ [14].

In 2006 [4], M.Jibladze and T. Pirashvili presented the definition of *categorical rings* as a slightly modified version of the definition of Ann-categories and classified them by Mac Lane ring cohomology. However, in [10] authors have showed that, it has not been proved whether the R -bimodule structure on M can be deduced from the axiomatics of categorical rings.

The Ann-functor classification problem has been solved for regular Ann-categories with Shukla cohomology [1, 12]. In this paper, we present a solution for this problem in the general case via low-dimensioned cohomology groups of Mac Lane ring cohomology. In proper, Hochschild algebra cohomology used to classification of strong Ann-functor.

In this paper, for convenience, sometimes we denote by XY the tensor product of the two objects X and Y , instead of $X \otimes Y$.

2 Preliminaries

2.1 The basic concepts

The definition of Ann-categories was presented in [7, 10, 11]. We always suppose that $\mathcal{A}$ is an Ann-category with a system of constraints:

$$(a^+, c, (O, g, d), a, (I, l, r), \mathcal{L}, \mathcal{R}).$$

Definition 1. Let $\mathcal{A}$ and $\mathcal{A}'$ be Ann-categories. An Ann-functor from $\mathcal{A}$ to $\mathcal{A}'$ is a triple $(F, \check{F}, \tilde{F})$, where $(F, \check{F})$ is a symmetric monoidal functor respect to the operation $\oplus$, $(F, \tilde{F})$ is an A -functor (i.e. an associativity functor) respect to the operation $\otimes$, satisfies the two following commutative diagrams:

$$\begin{array}{ccccc} F(X(Y \oplus Z)) & \xleftarrow{\tilde{F}} & FX.F(Y \oplus Z) & \xleftarrow{id \otimes \tilde{F}} & FX(FY \oplus FZ) \\ \downarrow F(\Sigma) & & & & \downarrow \Sigma' \\ F(XY \oplus XZ) & \xleftarrow{\tilde{F}} & F(XY) \oplus F(XZ) & \xleftarrow{\tilde{F} \oplus \tilde{F}} & FX.FY \oplus FX.FZ \\ & & & & \downarrow \mathfrak{L}' \\ F((X \oplus Y)Z) & \xleftarrow{\tilde{F}} & F(X \oplus Y).FZ & \xleftarrow{\tilde{F} \otimes id} & (FX \oplus FY).FZ \\ \downarrow F(\mathfrak{R}) & & & & \downarrow \mathfrak{R}' \\ F(XZ \oplus YZ) & \xleftarrow{\tilde{F}} & F(XZ) \oplus F(YZ) & \xleftarrow{\tilde{F} \oplus \tilde{F}} & FX.FZ \oplus FY.FZ \end{array}$$

The commutation of the above diagrams are called the compatibility of the functor F with the distributivity constraints of the two Ann-categories $\mathcal{A}, \mathcal{A}'$.

We call $\varphi : F \rightarrow G$ an Ann-morphism between two Ann-functors $(F, \check{F}, \tilde{F})$ and $(G, \check{G}, \tilde{G})$ if it is an $\oplus$ -morphism as well as an $\otimes$ -morphism.

An Ann-functor $(F, \check{F}, \tilde{F}) : \mathcal{A} \rightarrow \mathcal{A}'$ is called an Ann-equivalence if there exists an Ann-functor $(F', \check{F}', \tilde{F}') : \mathcal{A}' \rightarrow \mathcal{A}$ and natural isomorphisms $\alpha : F \circ F' \cong id_{\mathcal{A}'}$, $\alpha' : F' \circ F \cong id_{\mathcal{A}}$.

By Theorem 8 [9], an Ann-functor $(F, \check{F}, \tilde{F}) : \mathcal{A} \rightarrow \mathcal{A}'$ is an Ann-equivalence iff F is a categorical equivalence.

Note that, similar to a ring homomorphism, an Ann-functor F is not required $F(1) \cong 1'$. Moreover, note that: in the Definition 1, it is only required that $(F, \tilde{F})$ is an AC -functor (i.e. an $\oplus$ -functor which is compatible with the associativity and commutativity constraints). Indeed, since $(\mathcal{A}, \oplus), (\mathcal{A}', \oplus)$ are Gr -categories, each A -functor is compatible with the unitivity constraints.

2.2 The third Mac Lane ring cohomology group $H_{MaL}^3(R, M)$

Let R be a ring and M be an R -bimodule. From the definition of Mac Lane ring cohomology [6], we may obtain the description of the elements of cohomology group $H_{MaL}^3(R, M)$.

The group $Z_{MaL}^3(R, M)$ of 3-cochains of the ring R , with coefficients in R -bimodules M , consisting of quadruples $(\sigma, \alpha, \lambda, \rho)$, functions:

$$\alpha, \lambda, \rho : R^3 \rightarrow M$$

and $\sigma : R^4 \rightarrow M$ satisfy the following relations:

$$\text{M1. } x\alpha(y, z, t) - \alpha(xy, z, t) + \alpha(x, yz, t) - \alpha(x, y, zt) + \alpha(x, y, z)t = 0$$

$$\text{M2. } \alpha(x, z, t) + \alpha(y, z, t) - \alpha(x + y, z, t) + \rho(xz, yz, t) - \rho(x, y, zt) + \rho(x, y, z)t = 0$$

- M3. $-\alpha(x, y, t) - \alpha(x, z, t) + \alpha(x, y + z, t) + x\rho(y, z, t) - \rho(xy, xz, t) - \lambda(x, yt, zt) + \lambda(x, y, z)t = 0$
- M4. $\alpha(x, y, z) + \alpha(x, y, t) - \alpha(x, y, z + t) + x\lambda(y, z, t) - \lambda(xy, z, t) + \lambda(x, yz, yt) = 0$
- M5. $\lambda(x, z, t) + \lambda(y, z, t) - \lambda(x + y, z, t) + \rho(x, y, z) + \rho(x, y, t) - \rho(x, y, z + t) + \sigma(xz, xt, yz, yt) = 0$
- M6. $\lambda(x, a, b) + \lambda(x, c, d) - \lambda(x, a + c, b + d) - \lambda(x, a, c) - \lambda(x, b, d) + \lambda(x, a + b, c + d) - x\sigma(a, b, c, d) + \sigma(xa, xb, xc, xd) = 0$
- M7. $\rho(a, b, x) + \rho(c, d, x) - \rho(a + c, b + d, x) - \rho(a, c, x) - \rho(b, d, x) + \rho(a + b, c + d, x) - \sigma(ax, bx, cx, dx) + \sigma(a, b, c, d)x = 0$
- M8. $\sigma(a, b, c, d) + \sigma(x, y, z, t) - \sigma(a + x, b + y, c + z, d + t) + \sigma(a, b, x, y) + \sigma(c, d, z, t) - \sigma(a + c, b + d, x + z, y + t) + \sigma(a, c, x, z) + \sigma(b, d, y, t) - \sigma(a + b, c + d, x + y, z + t) = 0$
- M9. $\alpha(0, y, z) = \alpha(x, 0, z) = \alpha(x, y, 0) = 0$
- M10. $\sigma(0, 0, z, t) = \sigma(x, y, 0, 0) = \sigma(0, y, 0, t) = \sigma(x, 0, z, 0) = \sigma(x, 0, 0, t) = 0$.

The subgroup $B_{MaL}^3(R, M) \subset Z_{MaL}^3(R, M)$ of 3-coboundaries consists of the quadruples $(\sigma, \alpha, \lambda, \rho)$ such that there exist the maps $\mu, \nu : R^2 \rightarrow M$ satisfying:

- M11. $\sigma(x, y, z, t) = -\mu(x, y) - \mu(z, t) + \mu(x + z, y + t) + \mu(x, z) + \mu(y, t) - \mu(x + y, z + t)$
- M12. $\alpha(x, y, z) = x\nu(y, z) - \nu(xy, z) + \nu(x, yz) - \nu(x, y)z$
- M13. $\lambda(x, y, z) = \nu(x, y) + \nu(x, z) - \nu(x, y + z) + x\mu(y, z) - \mu(xy, xz)$
- M14. $\rho(x, y, z) = -\nu(x, z) - \nu(y, z) + \nu(x + y, z) + \mu(xz, yz) - \mu(x, y)z$.

Finally, $H_{MaL}^3(R, M) = Z_{MaL}^3(R, M) / B_{MaL}^3(R, M)$.

Each Ann-category $\mathcal{I}$ of the type (R, M) having the *structure* f is a family $f = (\xi, \eta, \alpha, \lambda, \rho)$, where $\xi, \alpha, \lambda, \rho : R^3 \rightarrow M$ and $\eta : R^2 \rightarrow M$ are functions satisfying 17 the equations (see Proposition 5.8 [11]). Now, we define a function $\sigma : R^4 \rightarrow M$, given by:

$$\sigma(x, y, z, t) = \xi(x + y, z, t) - \xi(x, y, z) + \eta(y, z) + \xi(x, z, y) - \xi(x + z, y, t).$$

This function is respect to the associativity-commutativity constraint v in the Ann-category $\mathcal{A}$, where

$$v = v_{X,Y,Z,T} : (X \oplus Y) \oplus (Z \oplus T) \longrightarrow (X \oplus Z) \oplus (Y \oplus T)$$

is given by commutative diagram fowllowing:

$$\begin{array}{ccccc} (X \oplus Y) \oplus (Z \oplus T) & \xrightarrow{a_+} & ((X \oplus Y) \oplus Z) \oplus T & \xleftarrow{a_+ \oplus T} & (X \oplus (Y \oplus Z)) \oplus T \\ \downarrow v & & & & \downarrow (X \oplus c) \oplus T \\ (X \oplus Z) \oplus (Y \oplus T) & \xrightarrow{a_+} & ((X \oplus Z) \oplus Y) \oplus T & \xleftarrow{a_+ \oplus T} & (X \oplus (Z \oplus Y)) \oplus T \end{array}$$

The quadruple $h = (\sigma, \alpha, \lambda, \rho)$ is a 3-cocycle of the ring R with coefficients in R -bimodule M due to Mac Lane (Theorem 7.2 [11]) and therefore each reduced Ann-category is of the form (R, M, h) .

3 An equivalence criterion of an Ann-functor

First, we show a characterized property of Ann-functors, which is related to the associativity-commutativity constraint v .

Definition 2. Let $\mathcal{A}, \mathcal{A}'$ be symmetric monoidal $\oplus$ -categories. Then, the $\oplus$ -functor $(F, \check{F}) : \mathcal{A} \rightarrow \mathcal{A}'$ is called compatible with the constraints v, v' if the following diagram commutes for all $X, Y, Z, T \in \mathcal{A}$

$$\begin{array}{ccccc}
 F((X \oplus Y) \oplus (Z \oplus T)) & \xleftarrow{\check{F}} & F(X \oplus Y) \oplus F(Z \oplus T) & \xleftarrow{\check{F} + \check{F}} & (FX \oplus FY) \oplus (FZ \oplus FT) \\
 \downarrow F(v) & & & & \downarrow v' \\
 F((X \oplus Z) \oplus (Y \oplus T)) & \xleftarrow{\check{F}} & F(X \oplus Z) \oplus F(Y \oplus T) & \xleftarrow{\check{F} + \check{F}} & (FX \oplus FZ) \oplus (FY \oplus FT)
 \end{array} \tag{1}$$

Then **Lemma 3.1.** Let $\oplus$ -functor $(F, \check{F}) : \mathcal{A} \rightarrow \mathcal{A}'$ be compatible with the unitivity constraints. Then $(F, \check{F})$ is an AC-functor iff it is compatible with the constraints v, v' .

Proof. The necessary condition was presented by D. B. A. Epstein (Lemma 1.5 [2]).

Now, assume that the diagram (1) commutes. To prove that the pair $(F, \check{F})$ is compatible with the commutativity constraints, we consider the following Diagram 1.

In the Diagram 1, the region (I) commutes thanks to the naturality of the morphism v , the regions (II) and (IV) commute since $(F, \check{F})$ is compatible with the unitivity constraints, the regions (III) and (VII) commute by the coherence theorem in a symmetric monoidal category, the regions (VI) and (VIII) commute thanks to the naturality of $\check{F}$, the outside region commutes by the diagram (1). Hence, the region (V) commutes. So $(F, \check{F})$ is compatible with the commutativity constraints.

$$\begin{array}{ccccc}
 F((O \oplus X) \oplus (Y \oplus O)) & \xrightarrow{F(v)} & F((O \oplus Y) \oplus (X \oplus O)) & & \\
 \uparrow \check{F} & \searrow F(g \oplus d) & \text{(VII)} & \swarrow F(g \oplus d) & \uparrow \check{F} \\
 & F(X \oplus Y) & \xrightarrow{F(c)} & F(Y \oplus X) & \\
 \text{(VI)} & & & & \text{(VIII)} \\
 F(O \oplus X) \oplus F(Y \oplus O) & \xrightarrow{\check{F}} & F(X \oplus Y) & \xrightarrow{\check{F}} & F(Y \oplus X) \\
 \uparrow \check{F} \oplus \check{F} & \searrow F(g) \oplus F(d) & \text{(V)} & \swarrow F(g) \oplus F(d) & \uparrow \check{F} \oplus \check{F} \\
 & FX \oplus FY & \xrightarrow{c'} & FY \oplus FX & \\
 \text{(II)} & \xrightarrow{g' \oplus d'} & \text{(III)} & \xrightarrow{g' \oplus d'} & \text{(IV)} \\
 (O \oplus FX) \oplus (FY \oplus O) & \xrightarrow{v'} & (O \oplus FY) \oplus (FX \oplus O) & & \\
 \uparrow (\hat{F} \oplus id) \oplus (id \oplus \hat{F}) & \text{(I)} & \uparrow (\hat{F} \oplus id) \oplus (id \oplus \hat{F}) & & \\
 (FO \oplus FX) \oplus (FY \oplus FO) & \xrightarrow{v'} & (FO \oplus FY) \oplus (FX \oplus FO) & &
 \end{array}$$

Diagram 1

Next, we consider the following Diagram 2.

In the Diagram 2, the region (I) commutes thanks to the naturality of the morphism v ; the first component of the region (II) commutes since $(F, \check{F})$ is compatible with unitivity constraints, the second one commutes thanks to the composition of morphisms, so the region (II) commutes; the regions (III) and (X) commute thanks to the coherence in a symmetric monoidal category; the first component of the region (IV) commutes thanks to the composition of morphisms, the second one commutes since $(F, \check{F})$ is compatible with unitivity constraints, so the region (IV) commutes; the region (V) and (VII) commute

thanks to the composition of morphisms; the regions (VIII) and (IX) commute thanks to the naturality of $\check{F}$; the outside region commutes thanks to the diagram (1). Therefore, the region (V) commutes, i.e., the pair $(F, \check{F})$ is compatible with associativity constraints.




Diagram 2

□

Proposition 3.2.

In the definition of Ann-functors, the condition that $(F, \check{F})$ is an symmetric monoidal $\oplus$ -functor is equivalent to the two following conditions:

1. $(F, \check{F})$ is compatible with the unitivity constraints respect to the operation $\oplus$,
2. $(F, \check{F})$ is compatible with the constraints v, v' .

Proof. Directly deduced from Lemma 3.1.

□

4 Ann-functors and the low-dimensioned cohomology groups of rings due to Mac Lane

4.1 Ann-functors of the type (p, q)

Now, we will show that each Ann-functor $(F, \check{F}, \tilde{F}) : \mathcal{A} \rightarrow \mathcal{A}'$ induces a Ann-functor $\overline{F}$ on their reduced Ann-categories, and this correspondence is 1-1. First, we have

Theorem 4.1. [Theorem 4.6 [11]] *Let $\mathcal{A}$ and $\mathcal{A}'$ be Ann-categories. Then, each Ann-functor $(F, \check{F}, \tilde{F}) : \mathcal{A} \rightarrow \mathcal{A}'$ induces the pair of ring homomorphisms:*

$$\begin{array}{ccc} F_0 : \Pi_0(\mathcal{A}) & \rightarrow & \Pi_0(\mathcal{A}') \\ \text{cls } X & \mapsto & \text{cls } FX \end{array} \quad ; \quad \begin{array}{ccc} F_1 : \Pi_1(\mathcal{A}) & \rightarrow & \Pi_1(\mathcal{A}') \\ u & \mapsto & \gamma_{F_0}^{-1}(Fu) \end{array}$$

satisfying

$$F_1(su) = F_0(s)F_1(u); \quad F_1(us) = F_1(u)F_0(s)$$

where $\Pi_1(\mathcal{A})$ is regarded as a ring with the null multiplication. Furthermore, F is a equivalence iff F_0, F_1 are isomorphisms.

The pair (F_0, F_1) is called the pair of induced homomorphisms of the Ann-functor $(F, \check{F}, \tilde{F})$. If $\mathcal{S}, \mathcal{S}'$ are, respectively, the reduced Ann-categories of $\mathcal{A}, \mathcal{A}'$ then the functor $\overline{F} : \mathcal{S} \rightarrow \mathcal{S}'$ given by

$$\overline{F}(s) = F_0(s), \quad \overline{F}(s, u) = (F_0 s, F_1 u)$$

is called the reduced functor of $(F, \check{F}, \tilde{F})$ on reduced Ann-categories.

Proposition 4.2. Let $\overline{F}$ be the induced functor of the Ann-functor $(F, \check{F}, \tilde{F}) : \mathcal{A} \rightarrow \mathcal{A}'$. Then the diagram

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{F} & \mathcal{A}' \\ H \uparrow & & \downarrow G' \\ \mathcal{S} & \xrightarrow{\overline{F}} & \mathcal{S}' \end{array}$$

commutes, where H, G' are canonical Ann-equivalences, and therefore $\overline{F}$ induces an Ann-functor.

Proof. This Proposition is naturally extended of Proposition 2 [8]. $\square$

Definition 3. Let $\mathcal{S} = (R, M, h)$, $\mathcal{S}' = (R', M', h')$ be Ann-categories. A functor $F : \mathcal{S} \rightarrow \mathcal{S}'$ is called a functor of the type (p, q) if

$$F(x) = p(x), \quad F(x, a) = (p(x), q(a)),$$

where $p : R \rightarrow R'$ is a ring homomorphism and $q : M \rightarrow M'$ is a group homomorphism satisfying

$$q(xa) = p(x)q(a), \quad q(ax) = q(a)p(x),$$

for $x \in R, a \in M$.

Proposition 4.3. Let $\mathcal{A} = (R, M, h)$, $\mathcal{A}' = (R', M', h')$ be Ann-categories and $(F, \check{F}, \tilde{F})$ is an Ann-functor from $\mathcal{A}$ to $\mathcal{A}'$. Then, $(F, \check{F}, \tilde{F})$ is a functor of the type (p, q) .

Proof. For $x, y \in R$, we have

$$\check{F}_{x,y} : F(x) \oplus F(y) \rightarrow F(x \oplus y), \quad \tilde{F}_{x,y} : F(x) \otimes F(y) \rightarrow F(x \otimes y)$$

are morphisms in the Ann-category $\mathcal{A}'$. Hence, $F(x) + F(y) = F(x + y)$ and $F(x) \cdot F(y) = F(xy)$, so the map $p : R \rightarrow R'$ given by $p(x) = F(x)$ is a ring homomorphism.

Assume that $F(x, a) = (p(x), q_x(a))$. Since $(F, \check{F})$ is a Gr-functor, according to Theorem 5 [8], $q_x = q$ for all $x \in R$. Moreover q is a group homomorphism:

$$q(a + b) = q(a) + q(b) \tag{2}$$

for all $a, b \in M$.

Since $(F, \tilde{F})$ is a $\otimes$ -functor, the following diagram

$$\begin{array}{ccc} Fx \otimes Fy & \xrightarrow{\tilde{F}} & F(x \otimes y) \\ F((x,a)) \otimes F((y,b)) \downarrow & & \downarrow F((x,a) \otimes (y,b)) \\ Fx \otimes Fy & \xrightarrow{\tilde{F}} & F(x \otimes y) \end{array}$$

commutes, for all morphisms $(x, a), (y, b)$. So, we have:

$$\begin{aligned} F((x, a) \otimes (y, b)) &= F(x, a) \otimes F(y, b) \\ \Leftrightarrow q_{xy}(ay + xb) &= q_x(a)F(y) + F(x)q_y(b) \end{aligned} \quad (3)$$

Applying $q_x = q_y = q_{xy} = q$ to the relation (3), we have:

$$q(ay + xb) = q(a)F(y) + F(x)q(b) \quad (4)$$

Applying $x = 1$ to (4), we have:

$$q(ay) = q(a)F(y) = q(a)p(y) \quad (5)$$

Applying $y = 1$ to (4), we have:

$$q(xb) = F(x)q(b) = p(x)q(b) \quad (6)$$

If R' -bimodule M' is regarded as an R -bimodule thanks to the actions $xa' = p(x)a', a'x = a'p(x)$, from the equations (2), (5), (6) we may show that $q : M \rightarrow M'$ is a homomorphism between R -bimodules. $\square$

4.2 Classification of Ann-functors

The existence problem of Ann-functors between Ann-categories has been solved for the regular Ann-categories (Theorem 5.1 [13], Theorem 4.2 [1]) thanks to Shukla cohomology. In this section, we will solve that problem in the general case.

Definition 4. If $F : (R, M, h) \rightarrow (R', M', h')$ is a functor of the type (p, q) , then F induces 3-cocycles $h_* = q_*h = q(h)$, $h'^* = p^*h' = h'p$, for example

$$\begin{aligned} \sigma'^*(x, y, z, t) &= \sigma'(p(x), p(y), p(z), p(t)) \\ \sigma_*(x, y, z, t) &= q(\sigma(x, y, z, t)). \end{aligned}$$

The function $k = p^*h' - q_*h$ is called an obstruction of the functor of the type (p, q) . Then we have

Theorem 4.4. The functor $F : (R, M, h) \rightarrow (R', M', h')$ of the type (p, q) is an Ann-functor iff the obstruction $\bar{k} = 0$ in $H_{MaL}^3(R, M')$.

Proof. Let $(F, \check{F}, \tilde{F}) : (R, M, h) \rightarrow (R', M', h')$ be an Ann-functor of the type (p, q) . Since $\check{F}_{x,y} = (\bullet, \mu(x, y))$, $\tilde{F}_{x,y} = (\bullet, \nu(x, y))$ where $\mu, \nu : R^2 \rightarrow M'$, we may identify $\check{F}, \tilde{F}$ with μ, ν and call μ, ν the pair of associated functions with $\check{F}, \tilde{F}$. According to Lemma 3.1, $(F, \check{F}, \tilde{F})$ is compatible with the pair of constraints (v, v') , i.e. the diagram (1) commutes, so we have:

$$\begin{aligned} 7. \quad \sigma'^*(x, y, z, t) - \sigma_*(x, y, z, t) &= \mu(x, y) + \mu(z, t) - \mu(x + z, y + t) - \mu(x, z) \\ &\quad - \mu(y, t) + \mu(x + y, z + t) \end{aligned}$$

Since F is compatible with the associativity constraint of multiplication, the distributivity constraints of Ann-categories $\mathcal{A}$ and $\mathcal{A}'$, we have:

$$8. \quad \alpha'^*(x, y, z) - \alpha_*(x, y, z) = x\nu(y, z) - \nu(xy, z) + \nu(x, yz) - \nu(x, y)z$$

$$9. \quad \lambda'^*(x, y, z) - \lambda_*(x, y, z) = \nu(x, y + z) - \nu(x, y) - \nu(x, z) + x\mu(y, z) - \mu(xy, xz)$$

$$10. \quad \rho'^*(x, y, z) - \rho_*(x, y, z) = \nu(x + y, z) - \nu(x, z) - \nu(y, z) + \mu(x, y)z - \mu(xz, yz)$$

From the equations (7)-(10), we have:

$$h'^* - h_* = \delta g \quad (11)$$

where $g = (-\mu, \nu)$. Hence the obstruction of the functor F vanishes in the cohomology group $H_{MaL}^3(R, M)$.

Conversely, assume that the obstruction of the functor F vanishes in the cohomology group $H_{MaL}^3(R, M')$. Then there exists a 2-cochain $g = (\mu, \nu)$ such that $h'^* - h_* = \delta g$. Take $\check{F}, \tilde{F}$ be functor morphisms associated with the functions $-\mu, \nu$, we can verify that $(F, \check{F}, \tilde{F})$ is an Ann-functor. $\square$

Theorem 4.5. *If there exists an Ann-functor $(F, \check{F}, \tilde{F}) : \mathcal{A} \rightarrow \mathcal{A}'$, of the type (p, q) then:*

1. *There exists a bijection between the set of congruence classes of Ann-functors of the type (p, q) and the cohomology group $H_{MaL}^2(R, M')$ of the ring R with coefficients in R -bimodule M' .*

2. *There exists a bijection*

$$\text{Aut}(F) \rightarrow Z_{MaL}^1(R, M')$$

between the group of automorphisms of the Ann-functor F and the group $Z_{MaL}^1(R, M')$.

Proof. (a) Suppose that there exists $(F, \check{F}, \tilde{F}) : \mathcal{A} \rightarrow \mathcal{A}'$, which is an Ann-functor of the type (p, q) . According to Theorem 4.4, we have

$$\overline{h'^* - h_*} = 0.$$

Hence, there exists a 2-cochain k such that

$$h'^* - h_* = \delta k.$$

Fix 2-cochain k . Now, we assume that

$$(G, \check{G}, \tilde{G}) : (R, M, h) \rightarrow (R', M', h')$$

is an Ann-functor of the type (p, q) . Then, from the proof of the Theorem 4.4, we have

$$h'^* - h_* = \delta g.$$

Hence, $k - g$ is a 2-cocycle. Consider the correspondence:

$$\Phi : \text{class}(G) \mapsto \text{class}(k - g)$$

from the set of the congruence classes of Ann-functors of the type (p, q) to the group $H_{MaL}^2(R, M')$.

First, we prove that the above correspondence is a map. Indeed, suppose that

$$(G', \check{G}', \tilde{G}') : (R, M, h) \rightarrow (R', M', h')$$

is also an Ann-functor of the type (p, q) and $u : G \rightarrow G'$ is an Ann-morphism. Since u is an $\oplus$ -morphism as well as an $\otimes$ -morphism, we have:

$$g' = g - \delta(u) \quad (12)$$

So

$$k - g' = k - g + \delta(u).$$

Thus $\overline{k - g} = \overline{k - g'} \in H_{MaL}^2(R, M')$.

Now, we prove that Φ is an injection. Assume that

$$(G, \check{G}, \tilde{G}), (G', \check{G}', \tilde{G}') : (R, M, h) \rightarrow (R', M', h')$$

are Ann-functors of the type (p, q) and satisfying

$$\overline{k - g} = \overline{k - g'} \in H_{MaL}^2(R, M').$$

Then, there exists an 1-cochain u such that

$$k - g = k - g' - \delta(u)$$

That means

$$g' = g - \delta(u).$$

Hence, the following diagrams:

$$\begin{array}{ccc} G(x) \oplus G(y) & \xrightarrow{\check{G}} & G(x \oplus y) \\ \downarrow u_x \oplus u_y & & \downarrow u_{x \oplus y} \\ G'(x) \oplus G'(y) & \xrightarrow{\check{G}'} & G'(x \oplus y) \end{array} \quad \begin{array}{ccc} G(x) \otimes G(y) & \xrightarrow{\tilde{G}} & G(x \otimes y) \\ \downarrow u_x \otimes u_y & & \downarrow u_{x \otimes y} \\ G'(x) \otimes G'(y) & \xrightarrow{\tilde{G}'} & G'(x \otimes y) \end{array}$$

commute, it means that $u : G \rightarrow G'$ is an Ann-morphism. Therefore,

$$class(G) = class(G').$$

Finally, we must prove that the correspondence Φ is a surjection. Indeed, assume that g is an arbitrary 2-cocycle. We have:

$$\delta(k - g) = \delta k - \delta g = \delta k = h'^* - h_*.$$

Then, according to Theorem 4.4, there exists an Ann-functor

$$(G, \check{G}, \tilde{G}) : (R, M, h) \rightarrow (R', M', h')$$

of the type (p, q) , and the pair of isomorphisms $\check{G}, \tilde{G}$ associated with the 2-cochain $k - g$.

Clearly, $\Phi(G) = \overline{g}$. So Φ is a surjection.

(b) Assume that $F = (F, \check{F}, \tilde{F}) : (R, M, h) \rightarrow (R', M', h')$ is an Ann-functor of the type (p, q) and $u \in Aut(F)$. Then, from the equation (12) with $g' = g$, we have $\delta(u) = 0$, i.e., $u \in Z_{MaL}^1(R, M')$. $\square$

5 Ann-functors and Hochschild cohomology

In this section, we will consider special Ann-functors which are related to the low-dimensional Hochschild groups.

Following, we will find a condition for the existence of Ann-functors of the form

$$F = (F, id, \tilde{F}) : (R, M, h) \rightarrow (R', M', h')$$

of the type $(p, 0)$, where $p : R \rightarrow R'$ is a ring homomorphism.

Suppose that there exists an Ann-functor

$$F = (F, id, \tilde{F} = \nu) : (R, M, h) \rightarrow (R', M', h')$$

of the type $(p, 0)$. Then, the equations (7) - (10) turn into:

13. $\sigma'^*(x, y, z, t) = 0$
14. $\alpha'^*(x, y, z) = x\nu(y, z) - \nu(xy, z) + \nu(x, yz) - \nu(x, y)z$
15. $\lambda'^*(x, y, z) = \nu(x, y + z) - \nu(x, y) - \nu(x, z)$
16. $\rho'^*(x, y, z) = \nu(x + y, z) - \nu(x, z) - \nu(y, z)$

and the Theorem 4.4 turns into

Corollary 5.1. *Let $p : R \rightarrow R'$ be a ring homomorphism. There exists an Ann-functor $(F, id, \tilde{F})$ from (R, M, h) to (R', M', h') of the type $(p, 0)$ iff $\overline{h'^*} = 0 \in H_{MaL}^3(R, M')$.*

Each cocycle of $\mathbb{Z}$ -algebras due to Hochschild is a multi-linear function. This suggests us the following definition:

Definition 5. *An Ann-functor*

$$(F, id, \tilde{F}) : (R, M, h) \rightarrow (R', M', h')$$

*of the type $(p, 0)$ is called a **strong** Ann-functor if the function $\nu : R^2 \rightarrow M'$ corresponding to $\tilde{F}$ is bi-additive.*

If ν is a normal bi-additive function then ν is a 2-cocycle of the $\mathbb{Z}$ -algebra R with coefficients in R -bimodule M' due to Hochschild. Then, in the equations (13)-(16), α'^* is a normal multi-linear function and other functions are equal to 0. So, we can identify

$$h'^* \equiv \alpha'^* = \delta(\nu),$$

in which $\delta(\nu)$ is an 3-coboundary of the ring R with coefficients in R -bimodule M' due to Hochschild. Then, we have the following proposition, as a direct corollary of Theorem 4.4.

Proposition 5.2. *Let $F : (R, M, h) \rightarrow (R', M', h')$ be a functor of the type $(p, 0)$. There exists a strong Ann-functor $(F, id, \tilde{F})$ iff its cohomology class $\overline{h'^*} = 0$ in the cohomology group $H_{Hochs}^3(R, M')$.*

Theorem 5.3. *If there exists an strong Ann-functor $(F, id, \tilde{F}) : (R, M, h) \rightarrow (R', M', h')$, of the type $(p, 0)$, then:*

1. *There exists a bijection between the set of congruence classes of strong Ann-functors of the type $(p, 0)$ and the cohomology group $H_{Hochs}^2(R, M')$ of the ring R with coefficients in R -bimodule M' .*
2. *There exists a bijection*

$$Aut(F) \rightarrow Z_{Hochs}^1(R, M')$$

between the group of automorphisms of the Ann-functor F and the group $Z_{Hochs}^1(R, M')$.

Proof. (a) The restriction Φ_H of the map Φ , referred in Theorem 4.5, on the set of congruence classes of strong Ann-functors, gives us an injection to the group $H_{Hochs}^2(R, M')$. Moreover, it is easy to see that Φ_H is also a surjection.

(b) Assume that $F = (F, id, \tilde{F}) : (R, M, h) \rightarrow (R', M', h')$ is an strong *Ann*-functor of the type $(p, 0)$ and $u \in Aut(F)$. Then u is bi-linear respect to the $\oplus$. So $u \in Z_{Hochs}^1(R, M')$. The converse also holds. $\square$

References

- [1] T. P. Dung, *Doctoral dissertation*, Hanoi, Vietnamese, 1992.
- [2] D. B. A. Epstein, *Functors between tenxored categories*, Invent. math.1, 221 -228 (1966).
- [3] G. Hochschild, *On the cohomology groups of an associative algebra*, Ann. of Math. (2) 46, (1945). 58-67.
- [4] M. Jibladze and T. Pirashvili, *Third Mac Lane cohomology via categorical rings*, arxiv. math. KT/0608519 v1, 21 Aug 2006.
- [5] S. Mac Lane, *Extensions and Obstruction for rings*, Illinois J. Mathematics, 2 (1958), 316-345.
- [6] S. Mac Lane, *Homologie des anneaux et des modules*, Colloque de Topologie algebrique, Louvain (1956), 55-88.
- [7] N. T. Quang, *Doctoral dissertation*, Hanoi, Vietnamese, 1988.
- [8] N. T. Quang, *On Gr-functors between Gr-categories: Obstruction theory for Gr-functors of the type (φ, f)* , arXiv: 0708.1348 v2 [math.CT] 18 Apr 2009
- [9] N. T. Quang and P. L. Hong Anh, *On monoidal equavalences and Ann-equavalences*, arXiv: 0705.0736 v1 [math. CT] 5 May 2007.
- [10] N. T. Quang, D. D. Hanh and N. T. Thuy, *On the axiomatics of Ann-categories*, JP Journal of Algebra, Number Theory and applications, Vol 11, No 1, 2008, 59 - 72.
- [11] N. T. Quang, *Structure of Ann-categories*, arXiv. 0805. 1505 v3 [math. CT] 6 Apr 2009.
- [12] N. T. Quang, *Ann-categories and the Mac Lane-Shukla cohomology of rings*, Abelian groups and modules No 11,12 (Russian), 166 - 183, Tomsk. Gos. Univ., Tomsk, 1994 .
- [13] N. T. Quang, *Structure of Ann-categories and Mac Lane-Shukla cohomology*, East-West, J. of Math. 5 (2003), 51-66.
- [14] U. Shukla, *Cohomologic des algbres associativ*, Ann. Sci. cole Norm. Sup. (3) 78 (1961), 163 - 209.

Dept. of Mathematics, Hanoi National University of Education, Viet Nam
Email: nguyenguang272002@gmail.com
ddhanhdhsphn@gmail.com